

Challenges and opportunities for the Bolivian Biodiversity Observation Network

Miguel Fernández, Laetitia M. Navarro, Amira Apaza-Quevedo, Silvia C. Gallegos, Alexandra Marques, Carlos Zambrana-Torrel, Florian Wolf, Healy Hamilton, Alvaro J. Aguilar-Kirigin, Luis F. Aguirre, Marcela Alvear, James Aparicio, Lilian Apaza-Vargas, Gabriel Arellano, Eric Armijo, Nataly Ascarrunz, Soraya Barrera, Stephan G. Beck, Héctor Cabrera-Condorco, Consuelo Campos-Villanueva, Leslie Cayola, N. Paola Flores-Saldana, Alfredo F. Fuentes, M. Carolina García-Lino, M. Isabel Gómez, Yara S. Higuera, Michael Kessler, Juan Carlos Ledezma, J. Miguel Limachi, Ramiro P. López, M. Isabel Loza, Manuel J. Macía, Rosa I. Meneses, Tatiana B. Miranda, A. Bruno Miranda-Calle, R. Fernando Molina-Rodríguez, Mónica Moraes R., M. Isabel Moya-Díaz, Mauricio Ocampo, Humberto L. Perotto-Baldivieso, Oscar Plata, Steffen Reichle, Kathia Rivero, Renate Seidel, Liliana Soria, Marcos F. Terán, Marisol Toledo, F. Santiago Zenteno-Ruiz & Henrique Miguel Pereira

To cite this article: Miguel Fernández, Laetitia M. Navarro, Amira Apaza-Quevedo, Silvia C. Gallegos, Alexandra Marques, Carlos Zambrana-Torrel, Florian Wolf, Healy Hamilton, Alvaro J. Aguilar-Kirigin, Luis F. Aguirre, Marcela Alvear, James Aparicio, Lilian Apaza-Vargas, Gabriel Arellano, Eric Armijo, Nataly Ascarrunz, Soraya Barrera, Stephan G. Beck, Héctor Cabrera-Condorco, Consuelo Campos-Villanueva, Leslie Cayola, N. Paola Flores-Saldana, Alfredo F. Fuentes, M. Carolina García-Lino, M. Isabel Gómez, Yara S. Higuera, Michael Kessler, Juan Carlos Ledezma, J. Miguel Limachi, Ramiro P. López, M. Isabel Loza, Manuel J. Macía, Rosa I. Meneses, Tatiana B. Miranda, A. Bruno Miranda-Calle, R. Fernando Molina-Rodríguez, Mónica Moraes R., M. Isabel Moya-Díaz, Mauricio Ocampo, Humberto L. Perotto-Baldivieso, Oscar Plata, Steffen Reichle, Kathia Rivero, Renate Seidel, Liliana Soria, Marcos F. Terán, Marisol Toledo, F. Santiago Zenteno-Ruiz & Henrique Miguel Pereira (2015) Challenges and opportunities for the Bolivian Biodiversity Observation Network, *Biodiversity*, 16:2-3, 86-98, DOI: [10.1080/14888386.2015.1068710](https://doi.org/10.1080/14888386.2015.1068710)

To link to this article: <http://dx.doi.org/10.1080/14888386.2015.1068710>




© 2015 The Author(s). Published by Taylor & Francis



Published online: 17 Aug 2015.



Submit your article to this journal [↗](#)

 Article views: 695

 [View related articles](#) 

 [View Crossmark data](#) 



Challenges and opportunities for the Bolivian Biodiversity Observation Network

Miguel Fernández^{a,b,c,*}, Laetitia M. Navarro^a, Amira Apaza-Quevedo^{a,d}, Silvia C. Gallegos^d, Alexandra Marques^a, Carlos Zambrana-Torrel^e, Florian Wolf^a, Healy Hamilton^f, Alvaro J. Aguilar-Kirigin^g, Luis F. Aguirre^h, Marcela Alvearⁱ, James Aparicio^g, Lilian Apaza-Vargas^j, Gabriel Arellano^k, Eric Armijo^l, Nataly Ascarrunz^m, Soraya Barrera^g, Stephan G. Beck^d, Héctor Cabrera-Condorcoⁿ, Consuelo Campos-Villanueva^d, Leslie Cayola^d, N. Paola Flores-Saldana^o, Alfredo F. Fuentes^d, M. Carolina García-Lino^p, M. Isabel Gómez^g, Yara S. Higuera^q, Michael Kessler^r, Juan Carlos Ledezma^s, J. Miguel Limachi^g, Ramiro P. López^d, M. Isabel Loza^t, Manuel J. Macía^k, Rosa I. Meneses^u, Tatiana B. Miranda^d, A. Bruno Miranda-Calle^g, R. Fernando Molina-Rodríguez^v, Mónica Moraes R.^d, M. Isabel Moya-Díaz^g, Mauricio Ocampo^g, Humberto L. Perotto-Baldivieso^w, Oscar Plata^d, Steffen Reichle^x, Kathia Rivero^{aa}, Renate Seidel^d, Liliana Soria^{aa}, Marcos F. Terán^y, Marisol Toledo^m, F. Santiago Zenteno-Ruiz^d and Henrique Miguel Pereira^{a,z}

^aGerman Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany;

^bDepartment of Integrative Biology, University of California, Berkeley, CA, USA;

^cInstituto de Ecología, Universidad Mayor de San Andrés, La Paz, Bolivia;

^dHerbario Nacional de Bolivia, Instituto de Ecología, Universidad Mayor de San Andrés, La Paz, Bolivia;

^eEcoHealth Alliance, New York, NY, USA;

^fNatureServe, Arlington, VA, USA;

^gColección Boliviana de Fauna, Museo Nacional de Historia Natural-Instituto de Ecología, La Paz, Bolivia;

^hCentro de Biodiversidad y Genética, Facultad de Ciencias y Tecnología, Universidad Mayor de San Simón, Cochabamba, Bolivia;

ⁱInstitute for Biodiversity Science and Sustainability, California Academy of Sciences, San Francisco, CA, USA;

^jFood and Agriculture Organization of the United Nations-Bolivia, La Paz, Bolivia;

^kDepartamento de Biología, Universidad Autónoma de Madrid, Madrid, Spain;

^lInvestigador Independiente, Santa Cruz, Bolivia;

^mInstituto Boliviano de Investigación Forestal, Universidad Autónoma Gabriel René Moreno, Santa Cruz, Bolivia;

ⁿServicio Nacional de Áreas Protegidas, Ministerio de Medio Ambiente y Agua, La Paz, Bolivia;

^oSociety for Conservation Biology – Bolivia Chapter, La Paz, Bolivia;

^pDepartamento de Botánica, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Concepción, Chile;

^qInstituto de Biodiversidad BIORENA, Universidad de San Francisco Xavier de Chuquisaca, Sucre, Bolivia;

^rInstitute of Systematic Botany, University of Zurich, Zurich, Switzerland;

^sConservación Internacional, La Paz, Bolivia;

^tDepartment of Biology, University of Missouri, St. Louis, MO, USA;

^uHerbario Nacional de Bolivia, Museo Nacional de Historia Natural, La Paz, Bolivia;

^vGEOBOLIVIA, Vicepresidencia del Estado Plurinacional de Bolivia, La Paz, Bolivia;

^wCaesar Kleberg Wildlife Research Institute, Texas A and M University–Kingsville, Kingsville, TX, USA;

^xInvestigador Independiente, Santa Cruz, Bolivia;

^yAsociación Boliviana para la Investigación y Conservación de Ecosistemas Andino Amazónicos, La Paz, Bolivia;

^zCIBIO/InBIO, Vairão, Portugal;

^{aa}Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma Gabriel Rene Moreno, Santa Cruz, Bolivia

(Received 23 April 2015; final version received 30 June 2015)

Pragmatic methods to assess the status of biodiversity at multiple scales are required to support conservation decision-making. At the intersection of several major biogeographic zones, Bolivia has extraordinary potential to develop a monitoring strategy aligned with the objectives of the Group on Earth Observations Biodiversity Observation Network (GEO BON). Bolivia, a GEO Observer since 2005, is already working on the adequacy of national earth observations towards the objectives of the Global Earth Observation System of Systems (GEOSS). However, biodiversity is still an underrepresented component in this initiative. The integration of biodiversity into Bolivia's GEO framework would confirm the need for a country level biodiversity monitoring strategy, fundamental to assess the progress towards the 2020 Aichi targets. Here we analyse and discuss two aspects of the process of developing such a strategy: (1) identification of taxonomic, temporal and spatial coverage of biodiversity data to detect both availability and gaps; and (2) evaluation of issues related to the acquisition, integration and analyses of multi-scale and multi-temporal biodiversity datasets. Our efforts resulted in the most comprehensive biodiversity database for the country of Bolivia, containing 648,534 records

*Corresponding author. Email: miguel.fernandez.trigoso@gmail.com

for 27,534 species referenced in time and space that account for 92.5% of the species previously reported for the country. We capitalise this information into recommendations for the implementation of the Bolivian Biodiversity Observation Network that will help ensure that biodiversity is sustained as the country continues on its path of development.

Keywords: Bolivia; biodiversity; big data integration; monitoring; baseline; GEO BON

Introduction

In the last two decades, aided by increased connectivity and easy access to data capturing devices and analytical tools, societies are witnessing a change in the paradigm on how to deal with information. We are rapidly moving from data control schemes to more collaboration, integration and sharing (Wallis, Rolando, and Borgman 2013). Within the biodiversity community, there is a plethora of initiatives dedicated to collecting data on multiple dimensions of biodiversity and at different spatial scales and resolutions such as the Living Planet Index (Collen et al. 2009), the Map of Life (Jetz, McPherson, and Guralnick 2012) and the PREDICTS database (Hudson et al. 2014). More data could lead to improved knowledge on how biodiversity is distributed in space and how it is changing over time; both components are essential for better informed policy-making and more accurate scenarios for conservation and management (Pereira et al. 2010; Schmeller et al. 2015).

However, quantity does not mean quality (Maldonado et al. 2015). One of the main problems is that not all biodiversity data was collected using a sampling design appropriate for monitoring in space and time. For instance, in the last global biodiversity assessment (Tittensor et al. 2014) only 55 out of 163 potential indicators were selected to measure progress towards conservation targets. Spatial coverage and availability of temporal series were amongst five criteria used to determine the indicator's suitability to measure change and/or response to change (Tittensor et al. 2014). Thus urgent harmonisation and standardisation of methods, protocols and quality control measures are needed (Pereira et al. 2013). However, while ontological alignment (i.e. correspondence among concepts) is a priority in data rich regions, this is not as important in data deficient regions where a baseline for monitoring change might not even exist. Incidentally, data deficient regions also happen to occur not only in areas of high conservation value but also in areas where the highest degrees of degradation occur (Collen et al. 2008; Pereira, Navarro, and Martins 2012), making the strategies to define monitoring priorities even more urgent (Hardisty and Roberts 2013).

The country of Bolivia, at the heart of South America, is a perfect example where high levels of biodiversity, deficient information and high degrees of degradation overlap. Baseline information even for the most well-known and charismatic species data is lacking (Vié, Hilton-Taylor, and Stuart 2009). For example, from the

389 mammalian species described for Bolivia, 106 species have stable populations while 84 species are declining (Tarifa and Aguirre 2009). Yet, for the remaining 199 mammalian species, the status of their populations is unknown since there is simply not enough information to estimate or infer a trend (Peñaranda and Simonetti 2015). Similarly, for the 266 species of amphibians reported for Bolivia, available data on possible threats or declines is mostly anecdotal, and long-term well-funded programmes that monitor populations and putative declines are nonexistent (De la Riva and Reichle 2014).

One of the most pressing environmental concerns of Bolivia is deforestation for large-scale mechanised agriculture, small-scale agriculture and cattle ranching (Killeen et al. 2007; Müller et al. 2012). Although deforestation rates were considered moderate for decades, relative to other countries in the region, the situation has changed dramatically in the first decade of the twenty-first century. Independently of whether we look at the lower estimates (0.49%; FAO 2010) or the higher estimates of deforestation (0.66%; Cuéllar et al. 2012), Bolivia is currently placed in the top 10 list of countries with the highest annual rates of forest loss in the world (FAO 2010), with potentially dramatic consequences for biodiversity (Pinto-Ledezma and Rivero Mamani 2014). Thus, establishing monitoring schemes in, for instance, deforested and control areas would be an essential step towards the assessment of consequences of human-induced land-use change for biodiversity in the country.

To provide a baseline against which to measure biodiversity change in countries with similar co-occurring conditions to Bolivia, centralisation, systematisation, archiving and curation of biodiversity data is urgently needed. The Essential Biodiversity Variables framework concept proposed by Pereira et al. (2013) provides an attractive framework for the development of national and sub-national initiatives. These monitoring initiatives can provide the knowledge base to assess the targets for 2020 set by the Convention on Biological Diversity (CBD).

As the first step towards building a monitoring scheme for biodiversity in Bolivia, we evaluate two key aspects in establishing a baseline: first, the identification of taxonomic, temporal and spatial data availability to detect both data gaps and opportunities for long-term monitoring; and second, the evaluation of issues related to the acquisition, integration and analyses of multi-scale biodiversity datasets. We capitalise this information into recommendations for the implementation of the Bolivian Biodiversity

Observation Network, consistent with the essential biodiversity variables framework.

Methods

Biodiversity observation: data acquisition and integration

We evaluated the taxonomic, spatial and temporal coverage of biodiversity observations in Bolivia from the year 1789 until 2015. We used, as the main sources of information on species occurrence, data derived from specimens hosted in natural history museums and herbaria located inside and outside Bolivia (i.e. Herbario Nacional de Bolivia (LPB) and Colección Boliviana de Fauna (CBF) – Universidad Mayor de San Andres, Museo de Historia Natural Noel Kempff Mercado (MHNNKM) – Universidad Autónoma Gabriel Rene Moreno, Missouri Botanical Garden (MOBOT), California Academy of Sciences (CAS), Museum of Vertebrate Zoology at Berkeley (MVZ), Smithsonian Institution National Museum of Natural History (NMNH), Museum National d'Histoire Naturelle de Paris (MNHN), Global Biodiversity Information Facility (GBIF)); we also included expert observations databases from other research institutions (e.g. Instituto Boliviano de Investigación Forestal (IBIF), Wildlife Conservation Society (WCS) and NatureServe). Biodiversity observations that fulfilled the following criteria: (1) terrestrial macrobiotic organisms identified at the species level, (2) with georeferenced occurrence information precise to minutes in latitude and longitude and (3) that contained information about the date of the collection event, were integrated into a single database using PostgreSQL, an open-source database software.

Once all the records were integrated into the database, we applied a series of consistent quality control routines on the data. These included a verification of the types of input values and alignment of fields to controlled vocabularies and standards (i.e. Darwin Core; Wiczorek et al. 2012). We also checked the taxonomic accuracy and redundancy of the database by aligning the scientific identification of each record against a reference taxonomic backbone. For plants we used the Taxonomic Name Resolution Service v.3.2. (TNRs; Boyle et al. 2013) and for animals and fungi we used the Integrated Taxonomic Information System database (ITIS 2010). After applying this systematic quality checks, we replaced synonymies by most current taxonomy and removed the invalid records from the database. We then transformed all the geographic coordinates to decimal degrees and using PostGIS, a spatial database extender, and we imported the database into a Geographic Information System (ArcGIS v.10.2).

Then, we checked if the coordinates of each record were aligned with the country (IDE-EPB 2015); we applied a buffer to the country boundaries and removed

all records that fell at distances larger than 2.5 km of the official international boundary of the country.

Taxonomic data coverage

From the integrated database, we first counted the number of species in taxonomic groups from which we were able to find previously published accounts (i.e. fish, amphibians, reptiles, birds, mammals, bryophytes and vascular plants) and then compared the results against our numbers. Secondly, using a national ecoregional classification based on Ibisich and Merida (2013), which provides a more tailored and accurate representation for Bolivia than the global classification from Olson and Dinerstein (2002), we counted the number of records and species that occur within each ecoregion. Based on the results of these analyses we calculated the ratio of the number of records to the number of species, and reported averaged values per ecoregion for animals and plants. A ratio close to one is an indicator that more species in the group are known from one single record.

Spatial and temporal data coverage

We created a grid of $5 \times 5 \text{ km}^2$, and counted the number of unique species that occur in each cell. We defined the spatial resolution of the grid based on the average error estimate of the retrospectively georeferenced observations (~2.5 km) using the point-radius method developed by Wiczorek, Guo, and Hijmans (2004), a well known and accepted method that has been incorporated in protocols, geospatial guidelines and software (e.g. BioGeomancer, SpeciesGeoCoder, ModEco). We also counted the number of unique years represented in each 25 km^2 cell, to understand the temporal distribution of the data across space. Finally, to evaluate the distribution of records across time in our database, we plotted the number of records and the number of species collected for each year. In order to compare national and regional patterns, we repeated these exercises at the ecoregion level.

Results

From a total of 965,896 biodiversity observations, our data integration efforts resulted in a database referenced at the taxonomic, spatial and temporal level that contains 648,534 records from 27,534 species for the country of Bolivia (Figure 1). From this database, 93.7% of the records were obtained from vouchers specimens from natural history museums and herbaria and the remaining 6.3% of the records were originated from direct expert observations. Also, 55.4% of the records were obtained from institutions hosted outside Bolivia and 44.6% from institutions hosted inside the country.

Our database contains 92.5% of the total number of species reported for Bolivia when compared with




Figure 1. Georeferenced biodiversity observations for Bolivia, compiled from records in local and international natural history museums, herbaria and direct observations reported by experts.

species accounts and checklists from previous studies (Figure 2) and 95.8% of the species included in the IUCN Red List of Threatened Species (Categories: CR, EN and VU), providing a good coverage of the number of species for the most well-known taxonomic groups including amphibians, birds and mammals. Our results also indicate that 34.5% of all the records fall within a protected area with an IUCN protection status, which indicates that biodiversity observations have been sampled slightly more inside protected areas given that 25% of the total area of the country is under protection.

Using ratios, we found that in average plants have 10 times less records per species than animals (Table 1). For animals, the ecoregion with the lowest number of records relative to the number of species was the *Prepuna*, with an average of two records per species; the larger number of records relative to the number of species were from *Sudoeste de la Amazonia* and *Gran Chaco* with an average of 21.3 and 22.7 records per species, respectively (Table 1). For plants, the ecoregions with the lowest number of records relative to species were *Lago Titicaca*, *Prepuna* and *Puna Sureña*, with an average of 1.6, 2.2 and 2.7 records per species,




Figure 2. Taxonomic representativity. Number of species from the integrated database (black) compared to number of species in previous studies (grey) from well-known taxonomic groups: fish (J. Sarmiento pers. com.), amphibians (De la Riva and Reichle 2014), reptiles (Aguirre, Aguayo, and Balderrama 2009), birds (S. Herzog pers. com.), mammals (Peñaranda and Simonetti 2015), bryophytes (Churchill, Sanjines-Asturizaga, and Aldana 2009) and vascular plants (Jørgensen, Nee, and Beck 2015).

Table 1. Number of records vs. number of species.

| Code | Ecoregion | Plants | | | Animals | | | Area sampled (%) |
|------|----------------------------|---------|---------|-------|---------|---------|-------|------------------|
| | | Records | Species | Ratio | Records | Species | Ratio | |
| PUSU | Puna Sur | 3092 | 1128 | 2.74 | 4645 | 312 | 14.89 | 8.54 |
| PREP | Prepuna | 660 | 289 | 2.28 | 108 | 53 | 2.04 | 12.3 |
| GRCH | Gran Chaco | 9682 | 3169 | 3.06 | 31,523 | 1388 | 22.71 | 15.37 |
| SAIN | Sabanas Inundables | 11,073 | 3335 | 3.32 | 34,110 | 2251 | 15.15 | 15.59 |
| BSCH | Bosque Seco Chiquitano | 11,630 | 3139 | 3.71 | 17,213 | 1306 | 13.18 | 15.61 |
| CERR | Cerrado | 14,310 | 4212 | 3.40 | 14,189 | 1352 | 10.49 | 15.68 |
| SAMZ | Sudoeste de la Amazonia | 126,373 | 9387 | 13.46 | 85,877 | 4026 | 21.33 | 22.24 |
| PUNO | Puna Norte | 21,088 | 4104 | 5.14 | 11,366 | 701 | 16.21 | 24.83 |
| LGTK | Lago Titicaca | 527 | 323 | 1.63 | 1650 | 168 | 9.82 | 27.89 |
| CHSE | Chaco Serrano | 7151 | 2222 | 3.22 | 9984 | 908 | 11.00 | 32.04 |
| BTBO | Bosque Tucumano Boliviano | 17,435 | 3821 | 4.56 | 9112 | 1077 | 8.46 | 39.65 |
| YUNG | Yungas | 112,198 | 10,466 | 10.69 | 33,915 | 2203 | 15.39 | 43.16 |
| BSIN | Bosques Secos Interandinos | 36,574 | 6639 | 5.51 | 23,049 | 1476 | 15.62 | 46.15 |

Notes: Number of records, species and the ratio between them, calculated per ecoregion for animals and plants. Ratio values larger than one indicate multiple records for a particular species.

respectively; and the largest number of records relative to species were from *Yungas* and *Sudoeste de la Amazonia*, with an average of 10 and 13 records per species, respectively (Table 1).

From a species richness perspective, at the resolution of 25 km², the number of species recorded per pixel ranged from 1 to 1278 with a mean of 27 species (Figure 3). The highest numbers of species were

recorded in *Yungas* (Figures 3 and 4). From a spatial perspective the ecoregions that were better sampled were: *Bosques Secos Interandinos* (46.1% of the total area sampled), *Yungas* (43.1%) and *Bosque Tucumano Boliviano* (39.6%; Table 1 and Figure 4). Conversely the ecoregions that were least sampled were *Puna Sureña* (8.5%), *Prepuna* (12.3%) and *Gran Chaco* (15.3%; Table 1 and Figure 4).




Figure 3. Number of unique species calculated based on a grid of $5 \times 5 \text{ km}^2$.

The number of non-consecutive years recorded in each pixel range from 1 to 90 with a mean value of 3 years (Figure 5). The ecoregions with more years recorded are: *Bosques Secos Interandinos* and *Sudoeste de la Amazonia* (Figure 5). The oldest record in our database was collected in the *Gran Chaco* ecoregion in the year 1789 (Figure 6). Ecoregions where collection efforts started early in time are *Gran Chaco*, *Bosque Seco Interandino* and *Sudoeste de la Amazonia*; whereas more recent collection efforts are from the *Prepuna* and *Chaco Serrano*. In general, for all ecoregions, the sampling effort increased in the late 1970s with a marked

peak in the number of records and species between 1995 and 2005 (Figure 6).

Discussion

When investigating biodiversity response to global change and human pressure, it is important to make a distinction between ‘biodiversity loss’ and ‘biodiversity alterations’ (Pereira, Navarro, and Martins 2012). Determining a baseline and monitoring changes are essential to distinguish the spatial scale of extinction (i.e. from local to global), to identify potential range shifts, and to




Figure 4. Spatial coverage of biodiversity observations per ecoregion based on a grid of $5 \times 5 \text{ km}^2$.

measure changes in communities. In this paper, we provide a state of the art regarding Bolivia's knowledge on biodiversity data and show how collaborative initiatives can help to overcome major resource limitations, and move forward with the creation of a baseline that can serve as the foundation for a biodiversity monitoring strategy in Bolivia.

Our results, by no means are complete. Specific taxonomic groups where our database falls short when compared with previous species accounts are fish, reptiles and bryophytes (Figure 2). As a consequence, this database will require to be constantly and dynamically updated as

new information becomes available if the goal is that it serves as a reference against which biodiversity loss and alterations could be monitored. Along these lines, taxonomically accurate and spatially well-distributed data collected at short time intervals are essential to produce reliable scenarios of biodiversity change (Fernandez 2013) that can be used to inform issues such as climate change, food security and public health.

Several limitations preclude biodiversity data integration in Bolivia. They can be grouped under four main non-exclusive categories: dispersion, availability, completeness and alignment. Data dispersion means that




Figure 5. Temporal data coverage of biodiversity observations, calculated based on the number of non-contiguous years represented in each $5 \times 5 \text{ km}^2$ cell.

information is dispersed across multiple researchers and research institutions, each with different data standards and data sharing policies. Data availability is related to how much of the biodiversity information is actually readily available for smooth integration into a database system. In most of the cases this is proportional to the amount of information that has been digitised in a museum or herbarium. However, not all the information available in digital form is complete. Old specimens without GPS coordinates are a good example of this, requiring additional work to retrospectively georeference the textual descriptions of the places where they were

collected. Finally, if the data is in one place, available and complete, it might still require considerable amount of ontological alignment due to issues that have to do with changes in the taxonomy, differences in database standards and different reference vocabularies.

Despite the limitations mentioned above, the present effort represents the most comprehensive biodiversity database for the country of Bolivia built with one specific goal in mind: the creation of a baseline that can provide an objective basis for directing future collection efforts that can serve to monitor change. With more than half million records, the present database represents a




Figure 6. Bolivian biodiversity observations over time. Top panel: total number of records and species over time. Bottom-left and -right panel: number of records and species, respectively, per ecoregion over time.

good approximation of the state of knowledge of biodiversity in Bolivia, corresponding to 92.5% of the species reported in the literature. In this sense, the high representativity of our results provides a good foundation to integrate, analyze and interpret the information in taxonomic, spatial and temporal dimensions.

Taxonomic dimension

Biodiversity research in Bolivia is unevenly distributed across the tree of life. Traditionally, the focus of biodiversity research has been on highly emblematic and charismatic groups such as mammals and birds (e.g. Butchart et al. 2005; Collen et al. 2009). Much less effort has been placed in other groups such as reptiles, amphibians, plants and insects; and the present database for Bolivia is not the exception. The main reason behind this is the lack of financial, technical and human resources enhanced by a worldwide downward trend in available funding for natural history assessments. Biodiversity institutions in Bolivia, despite some isolated efforts, do not have yet a working coordinated strategy and a stable source of funding that allows the long-term storage, maintenance, curation and expansion of biodiversity data holdings and inventories. Infrastructure and technology devoted to the collection and identification of specimens in general is either insufficient or extremely limited and with serious problems in maintenance in the best of the cases. Moreover there are almost no incentives for training local taxonomists in these underrepresented groups, which is worsened by the lack of employment opportunities for young biodiversity researchers inside the country. Finally the lack of a clear national strategy reverberates in the absence of an evaluation mechanism towards a national monitoring strategy that includes these taxonomically underrepresented groups (MMAyA 2014).

Spatial dimension

Current knowledge on biodiversity in Bolivia is spatially biased. With still vast unexplored regions, the scarcity in number of records for the: *Gran Chaco*, *Bosque Seco Chiquitano* and *Sabanas Inundables del Norte* ecoregions (Figure 4) might be attributed to the low accessibility and low human population density of these areas (Ibisch, Chive, et al. 2003; Larrea-Alcázar et al. 2011); for the *Puna Sureña* this might also be explained by the intrinsic low diversity of this ecoregion (Table 1). The relatively high number of records in time and space in the *Yungas* and *Bosques Secos Interandinos* ecoregions might be the result of the research focus that these two regions have received over the years (Ibisch, Gerkmann, et al. 2003). Particularly, the highly diverse *Yungas* ecoregion is also one of the most threatened areas in Bolivia (Kessler 2001), highlighting the importance and

opportunity to focus conservation as well as monitoring efforts in the area. Our results also indicate that the *Sucre de la Amazonia* and the *Yungas* ecoregions include high levels of biodiversity, which is in line with previous estimates where the two ecoregions and the transition between them are listed as high priority ecosystems due to the high species diversity and high number of endemics, respectively (Araujo et al. 2010; Moraes R. et al. 2014; Müller et al. 2003; Young 2007).

Temporal dimension

Data collection efforts in Bolivia have not been continuous and data shows that there has been a steady decline in the last decade. Our analysis revealed two periods of time where there was a considerable increase in the number of biodiversity observations collected per year. The first one corresponds to the establishment of the Instituto de Ecología in the late 1970s, a pioneer institution in the field of ecology and systematics in Bolivia (Baudoin and España 1997; Ibisch 2003a), followed closely by other research oriented universities and institutions. The second increase (between 1995 and 2005) coincides with the adoption of the CBD by Bolivia in 1994, and the endorsement of the National Strategy for the Conservation of Biodiversity in 2002 (MMAyA 2014). Also as a result of the international funding community focus on biodiversity and biodiversity research, this decade was characterised by high financial support from national and international institutions (FAN 2009). The observed decrease in collection efforts in the last decade might be attributed to two interlinked elements: first, the change of direction and focus in governmental policies from biodiversity to management; and second, the decrease in previously available funding opportunities for biodiversity research in the country.

Biodiversity as a key for development

Bolivia acknowledges the importance of biodiversity for its development. For example, in Bolivia's Poverty Reduction Strategy (IMF 2001), it is stated that 'biodiversity could come to represent an increase of about 10% in the Gross Domestic Product (GDP), if activities are developed in ethnic and ecotourism, mitigation of climate change and biodiversity services relating to biotechnology, ecological products, and others' (IMF 2001). More recently, the country has defined five major axis of action regarding biodiversity, including linking conservation with both human development and economic potential, while highlighting the need to establish, *inter alia*, 'legal, institutional, and political conditions to implement a sustainable model of biodiversity development' (MMAyA 2014). However, realising the full potential of biodiversity for development requires knowledge on the status and trends.

Bolivia identified several limitations to the conservation of biodiversity in its National Biodiversity Strategy and Action Plan (MDSP 2001), being the most important the ‘lack of scientific knowledge on natural regeneration, growth rates, population viabilities’, the ‘absence of a definition of priorities for scientific investigation resulting from a lack of coordination between academics for *in situ* and *ex situ* conservation’ and an ‘insufficient and deficient transfer of technology’ (MDSP 2001). Since then, and as shown in the results, the sampling and monitoring effort, and thus knowledge, have stalled with some exceptions: (1) the Madidi Project, which is a collaboration among the Missouri Botanical Garden (MBG) and the Herbario Nacional de Bolivia (LPB), since 2002 it has been monitoring 50 permanent plots of 1 ha in the *Yungas* ecoregion over an elevation gradient of ~3000 m; (2) the Instituto Boliviano de Investigación Forestal (IBIF), a network of permanent plots distributed across the *Sudoeste de la Amazonia* and *Bosque Seco Chiquitano* ecoregions also generating biodiversity information since 2002; and (3) the Global Observation Research Initiative in Alpine Environments (GLORIA) monitoring the *Puna* with permanent plots since 2006, along an elevational gradient of 1000 m.

To date, no harmonised observation system that delivers regular and timely data on biodiversity change exists in Bolivia that supports all levels of governance, management and decision-making. Despite the progress in biodiversity data integration and mobilisation ((e.g. Centro Geoespacial para la Biodiversidad de Bolivia (CGB; Perotto-Baldivieso et al. 2012)), Centro Digital de Recursos Naturales de Bolivia (CDRNB), it is still difficult for research institutions or even country level research infrastructures to develop, implement and maintain the platforms required to retrieve, share and leverage data investments through collaboration, integration and harmonisation. Only with the establishment of a well-funded national biodiversity monitoring strategy that can leverage individual efforts into a true collaboration and data sharing infrastructure, Bolivia will be able to take full advantage of the new data-intensive science that results from information integration needed to inform urgent pressing issues such as global change.

Acknowledgements

We are extremely grateful to all the Bolivian and foreign researchers funding agencies that contributed to the knowledge of biodiversity in Bolivia over the past fifty years; without their efforts this type of work would not have been possible.

Disclosure statement

All authors declare no conflict of interests.

Funding

MF, AAQ, LN, AM, FW and HMP were supported by the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig funded by the German Research Foundation (FZT 118). The rest of the co-authors were supported by their own institutions.

References

- Aguirre, L., R. Aguayo, and J. Balderrama. 2009. *Libro rojo de la fauna silvestre de vertebrados de Bolivia* [The Red Book of Vertebrates in Bolivia]. La Paz, Bolivia: Ministerio de Medio Ambiente y Agua.
- Araujo, N., R. Müller, C. Nowicki, and P. Ibsch. 2010. “Conclusiones y recomendaciones” [Conclusions and recommendations] In *Prioridades de la Conservación en Bolivia* [Conservation priorities in Bolivia], edited by N. Araujo, R. Müller, C. Nowicki, and P. Ibsch, 63–66. Bolivia: Fundación Amigos de la Naturaleza.
- Baudoin, M., and R. España. 1997. *Lineamientos para una estrategia nacional de conservación y uso sostenible de la diversidad biológica* [Guidelines for a National Strategy for Conservation and Sustainable Use of Biodiversity]. La Paz, Bolivia: MDSP/DGB/PNUD/UNOPS/TCA.
- Boyle, B., N. Hopkins, Z. Lu, J. Raygoza Garay, D. Mozzherin, T. Rees, N. Matasci, M. Narro, W. Piel, and S. McKay. 2013. “The Taxonomic Name Resolution Service: An Online Tool for Automated Standardization of Plant Names.” *BMC Bioinformatics* 14 (1): 16.
- Butchart, S., A. Stattersfield, J. Baillie, L. Bennun, S. Stuart, H. Akcakaya, C. Hilton-Taylor, and G. Mace. 2005. “Using Red List Indices to Measure Progress Towards the 2010 Target and Beyond.” *Philosophical Transactions of the Royal Society B: Biological Sciences* 360 (1454): 255–268.
- Churchill, S., N. Sanjines-Asturizaga, and C. Aldana. 2009. *Catálogo de las briofitas de Bolivia: diversidad, distribución y ecología* [Catalog of Bryophytes of Bolivia: Diversity, Distribution and Ecology]. Santa Cruz, Bolivia: La Rosa.
- Collen, B., M. Ram, T. Zamin, and L. McRae. 2008. “The Tropical Biodiversity Data Gap: Addressing Disparity in Global Monitoring.” *Tropical Conservation Science* 1 (2): 75–88.
- Collen, B., J. Loh, S. Whitmee, L. McRae, R. Amin, and J. Baillie. 2009. “Monitoring Change in Vertebrate Abundance: The Living Planet Index.” *Conservation Biology* 23 (2): 317–327.
- Cuéllar, S., A. Rodríguez, J. Arroyo, S. Espinoza, and D. Larrea. 2012. “Mapa de deforestación de las Tierras Bajas y los Yungas de Bolivia [Deforestation map of the Lowlands and Yungas of Bolivia] 2000–2005–2010”, edited by Fundación Amigos de la Naturaleza, p. 57. Santa Cruz, Bolivia.
- De la Riva, I., and S. Reichle. 2014. “Diversity and Conservation of the Amphibians of Bolivia.” *Herpetological Monographs* 28 (1): 46–65.
- FAN (Fundación Amigos de la Naturaleza). 2009. *20 Years Conserving Bolivia's Biodiversity, Memory*. Santa Cruz, Bolivia: Fundación Amigos de la Naturaleza.

- FAO (Food and Agriculture Organization of the United Nations). 2010. *Global Forest Resource Assessment*. Rome: Food and Agriculture Organization of the United Nations.
- Fernandez, M. 2013. "Consequences of Spatial and Temporal Climate Variability for Species Distribution Modeling." Ph.D., Environmental Engineering, University of California.
- Hardisty, A., and D. Roberts. 2013. "A Decadal View of Biodiversity Informatics: Challenges and Priorities." *BMC Ecology* 13 (1): 16.
- Hudson, L., T. Newbold, S. Contu, S. Hill, I. Lysenko, A. De Palma, H. Phillips, R. Senior, D. Bennett, and H. Booth. 2014. "The PREDICTS Database: A Global Database of How Local Terrestrial Biodiversity Responds to Human Impacts." *Ecology and Evolution* 4 (24): 4701–4735.
- Ibisch, P., and G. Merida. 2003. "Mapa de Ecoregiones de Bolivia" [Ecoregions map of Bolivia]. Fundacion Amigos de la Naturaleza FAN.
- Ibisch, P., J. Chive, S. Espinoza, and N. Araujo. 2003. "Hacia un mapa del estado de conservación de los ecosistemas de Bolivia" [Towards a map of the conservation status of the ecosystems in Bolivia]. In *Biodiversidad: La Riqueza de Bolivia* [Biodiversity: The Wealth of Bolivia], edited by P. Ibisch and G. Merida, 264–272. Santa Cruz, Bolivia: Fundación Amigos de la Naturaleza.
- Ibisch, P., B. Gerkmann, S. Kreft, S. Beck, S. Herzog, J. Kohler, R. Muller, S. Reichle, and R. Vásquez. 2003. "Consideraciones comparativas de patrones intercoregionales de diversidad de especies y de endemismo" [Comparative considerations in the inter-ecoregional patterns of species diversity and endemism]. In *Biodiversidad: La Riqueza de Bolivia* [Biodiversity: The Wealth of Bolivia], edited by P. Ibisch and G. Merida, 148–161. Santa Cruz, Bolivia: Fundación Amigos de la Naturaleza.
- IDE-EPB (Infraestructura de Datos Espaciales del Estado Plurinacional de Bolivia). 2015. "Infraestructura de Datos Espaciales del Estado Plurinacional de Bolivia" [Geo-spatial data infrastructure for the Plurinational State of Bolivia]. <http://geo.gob.bo/#map>.
- IMF (International Monetary Fund). 2001. *Poverty Reduction Strategy Paper*. Republic of Bolivia: Bolivian Government.
- ITIS (Integrated Taxonomic Information System). 2010. Integrated Taxonomic Information System On-Line Database. ITIS Partnership, T. Orrell, Acting Director, c/o Smithsonian Institution Washington, DC.
- Jetz, W., J. McPherson, and R. Guralnick. 2012. "Integrating Biodiversity Distribution Knowledge: Toward a Global Map Of Life." *Trends in Ecology & Evolution* 27 (3): 151–159.
- Jørgensen, P., M. Nee, and S. Beck. 2015. *Catálogo de las Plantas Vasculares de Bolivia* [Catalog of the vascular plants of Bolivia]. vol. 127. St. Louis, Missouri: Missouri Botanical Garden Press, St. Louis.
- Kessler, M. 2001. "Patterns of Diversity and Range Size of Selected Plant Groups Along An Elevational Transect in the Bolivian Andes." *Biodiversity & Conservation* 10 (11): 1897–1921.
- Killeen, T., V. Calderon, L. Soria, B. Quezada, M. Steininger, G. Harper, L. Solórzano, and C. Tucker. 2007. "'Thirty Years of Land-Cover Change in Bolivia'. AMBIO: A Journal of the Human." *Environment* 36 (7): 600–606.
- Larrea-Alcázar, D., D. Embert, L. Aguirre, B. Ríos-Uzeda, M. Quintanilla, and A. Vargas. 2011. "Spatial Patterns of Biological Diversity in a Neotropical Lowland Savanna of Northeastern Bolivia." *Biodiversity and Conservation* 20 (6): 1167–1182.
- Maldonado, C., C. I. Molina, A. Zizka, C. Persson, C. M. Taylor, J. Albán, E. Chilquillo, N. Rønsted, and A. Antonelli. 2015. "Estimating Species Diversity and Distribution in the Era of Big Data: To What Extent Can We Trust Public Databases?" *Global Ecology and Biogeography* 24: 973–984.
- MDSP (Ministerio de Desarrollo Sostenible y Planificación). 2001. *Estrategia Nacional de Conservación y Uso Sostenible de la Biodiversidad* [National Strategy for Conservation and Sustainable Use of Biodiversity]. La Paz, Bolivia: República de Bolivia.
- MMaA (Ministerio de Medio Ambiente y Agua). 2014. *Cuarto Informe Nacional convenio de las Naciones Unidas sobre a Diversidad Biológica* [4th National Report on the Convention of Biological Diversity]. La Paz, Bolivia: Estado Plurinacional de Bolivia.
- Moraes R., M., B. Ríos-Uzeda, L. Moreno, G. Huanca-Huarachi, and D. Larrea-Alcázar. 2014. "Using Potential Distribution Models for Patterns of Species Richness, Endemism, and Phytogeography of Palm Species in Bolivia." *Tropical Conservation Science* 7: 45–60.
- Müller, R., C. Nowicki, W. Barthlott, and P. Ibisch. 2003. "Biodiversity and Endemism Mapping as a Tool for Regional Conservation Planning—Case Study of the Pleurothallidinae (Orchidaceae) of the Andean Rain Forests in Bolivia." *Biodiversity & Conservation* 12 (10): 2005–2024.
- Müller, R., D. Müller, F. Schierhorn, G. Gerold, and P. Pacheco. 2012. "Proximate Causes of Deforestation in the Bolivian Lowlands: An Analysis of Spatial Dynamics." *Regional Environmental Change* 12 (3): 445–459.
- Olson, D., and E. Dinerstein. 2002. "The Global 200: Priority Ecoregions for Global Conservation." *Annals of the Missouri Botanical Garden* 89 (2): 199–224.
- Peñaranda, D., and J. Simonetti. 2015. "Predicting and Setting Conservation Priorities for Bolivian Mammals Based on Biological Correlates of the Risk of Decline." *Conservation Biology* 29 (3): 834–843.
- Pereira, H., P. Leadley, V. Proenca, R. Alkemade, J. Scharlemann, J. Fernandez-Manjarres, M. Araujo, P. Balvanera, R. Biggs, and W. Cheung. 2010. "Scenarios for Global Biodiversity in the 21st Century." *Science* 330 (6010): 1496–1501.
- Pereira, H., L. Navarro, and I. Martins. 2012. "Global Biodiversity Change: The Bad, the Good, and the Unknown." *Annual Review of Environment and Resources* 37: 25–50.
- Pereira, H., S. Ferrier, M. Walters, G. Geller, R. Jongman, R. Scholes, M. Bruford, N. Brummitt, S. Butchart, and A. Cardoso. 2013. "Essential Biodiversity Variables." *Science* 339 (6117): 277–278.
- Perotto-Baldivieso, H., K. Rivero, J. Pinto-Ledezma, and A. Gill. 2012. "Distributing Biodiversity Data Through the Web: The Geospatial Center for Biodiversity in Bolivia." Anais 4º Simpósio de Geotecnologias no Pantanal, Bonito, Brasil.

- Pinto-Ledezma, J., and M. Rivero Mamani. 2014. "Temporal Patterns of Deforestation and Fragmentation in Lowland Bolivia: Implications for Climate Change." *Climatic Change* 127 (1): 43–54.
- Schmeller, D., R. Julliard, P. Bellingham, M. Böhm, N. Brummitt, A. Chiarucci, D. Couvet, S. Elmendorf, D. Forsyth, and J. Moreno. 2015. "Towards a Global Terrestrial Species Monitoring Program." *Journal for Nature Conservation* 25: 51–57.
- Tarifa, T., and Aguirre L. F. 2009. "Mamíferos" [Mammals]. In *Libro rojo de la fauna de vertebrados de Bolivia* [The Red Book of Vertebrates in Bolivia], edited by Ministerio de Medio Ambiente y Agua, 419–572. La Paz, Bolivia: MMAA.
- Tittensor, D., M. Walpole, S. Hill, D. Boyce, G. Britten, N. Burgess, S. Butchart, P. Leadley, E. Regan, and R. Alkemade. 2014. "A Mid-term Analysis of Progress Toward International Biodiversity Targets." *Science* 346 (6206): 241–244.
- Vié, J., C. Hilton-Taylor, and S. Stuart. 2009. *Wildlife in a Changing World: An Analysis of the 2008 IUCN Red List of Threatened Species*. Gland, Switzerland: IUCN.
- Wallis, J., E. Rolando, and C. Borgman. 2013. "If We Share Data, Will Anyone Use Them? Data Sharing and Reuse in the Long Tail of Science and Technology." *PLoS ONE* 8 (7): e67332.
- Wieczorek, J., Qinghua Guo, and R. Hijmans. 2004. "The Point-Radius Method for Georeferencing Locality Descriptions and Calculating Associated Uncertainty." *International Journal of Geographical Information Science* 18 (8): 745–767.
- Wieczorek, J., D. Bloom, R. Guralnick, S. Blum, M. Döring, R. Giovanni, T. Robertson, and D. Vieglais. 2012. "Darwin Core: An Evolving Community-developed Biodiversity Data Standard." *PLoS ONE* 7 (1): e29715.
- Young, B. 2007. *Endemic Species Distributions on the East Slope of the Andes in Peru and Bolivia*. Arlington, VA: NatureServe.